

ST segment depression after Norwood/systemic-pulmonary artery shunt



Ashraf Harahsheh^{a,*}, Craig Sable^a, Pranava Sinha^b, Richard Jonas^b

^a Department of Pediatrics, Division of Cardiology, Children's National Health System, The George Washington University School of Medicine, Washington, DC

^b Department of Cardiovascular Surgery, Children's National Health System, The George Washington University School of Medicine, Washington, DC

^{a,b} USA

© 2014 King Saud University. Production and hosting by Elsevier B.V. All rights reserved.

Keywords: Diastolic runoff flow, Holter monitor, Single ventricle

A three-month-old girl with double inlet left ventricle (S/D/D), hypoplastic outlet ventricle, restrictive bulboventricular foramen, d-transposition of great arteries and interrupted aortic arch underwent Norwood stage 1 operation, systemic to pulmonary artery (PA) shunt and atrial septectomy, and developed ectopic atrial tachycardia that responded to digoxin therapy. She developed fussiness lasting two hours, not associated with cyanosis, and not relieved by feeding. Her pulse oximetry was 88% while breathing room air, and she had no differential blood pressure gradient. Apart from single second heart sound and a grade IV/VI ejection systolic murmur, the examination was normal. Her electrocardiogram and echocardiogram were unchanged. A Holter monitor revealed ST segment depression and T wave inversion upon increasing the heart rate beyond 140 beats/minute (Figs. 1 and 2). A cardiac catheterization showed patent systemic




Figure 1. Absence of ST segment depression with a heart rate <140 beats/minute in a patient with Norwood/systemic to pulmonary artery shunt with diastolic runoff flow.




Figure 2. ST segment depression and T wave inversion with a heart rate >140 beats/minute in a patient with Norwood/systemic to pulmonary artery shunt with diastolic runoff flow.

to PA shunt, patent native aorta, and no evidence of coarctation of aorta. After undergoing the second stage operation, her diastolic runoff

Disclosure: Authors have nothing to disclose with regard to commercial support.

Received 18 April 2014; accepted 29 April 2014.

Available online 20 May 2014

* Corresponding author. Address: Pediatric Cardiology, George Washington University, Center for Heart, Lung and Kidney Disease, Children's National Health System, 111 Michigan Ave, N.W. Washington, DC 20010, USA. Tel.: +202 476 2020; fax: +202 476 5700.

E-mail address: aharahsh@childrensnational.org (A. Harahsheh).



1016-7315 © 2014 King Saud University.

Production and hosting by Elsevier B.V. All rights reserved.

Peer review under responsibility of King Saud University.

URL: www.ksu.edu.sa

<http://dx.doi.org/10.1016/j.jsha.2014.04.007>



P.O. Box 2925 Riyadh - 11461KSA

Tel: +966 1 2520088 ext 40151

Fax: +966 1 2520718

Email: sha@sha.org.sa

URL: www.sha.org.sa



Production and hosting by Elsevier



Figure 3. Absence of ST depression with a heart rate <140 beats/minute after stage II palliation with no diastolic runoff flow.



Figure 4. Absence of ST depression with a heart rate >140 beats/minute after stage II palliation with no diastolic runoff flow.

and symptoms of fussiness disappeared and a surveillance Holter monitor showed better-looking ST segments and T waves even at higher heart rates (Figs. 3 and 4). Three months later, the patient is doing well and is awaiting Fontan completion.

Despite many advances in the field of pediatric cardiology, the interstage mortality rate remains high [1]. The use of the systemic to PA shunt in the Norwood operation has been complicated by wide pulse pressure, diastolic runoff flow [2,3], uneven distribution of pulmonary blood flow [4],

and increased early mortality [5]. Providers need to keep a high index of suspicion for the adverse effects of the diastolic runoff that accompanies the presence of systemic to PA shunts.

Funding sources or institutional or corporate affiliations

None.

References

- [1] Cross RR, Harahsheh AS, McCarter R, Martin GR. National pediatric cardiology quality improvement collaborative. identified mortality risk factors associated with presentation, initial hospitalisation, and interstage period for the Norwood operation in a multi-centre registry: a report from the national pediatric cardiology-quality improvement collaborative. *Cardiol Young* 2014;24(2): 253–62.
- [2] Ohye RG, Ludomirsky A, Devaney EJ, Bove EL. Comparison of right ventricle to pulmonary artery conduit and modified Blalock-Taussig shunt hemodynamics after the Norwood operation. *Ann Thorac Surg* 2004;78(3):1090–3.
- [3] Caspi J, Pettitt TW, Mulder T, Stopa A. Development of the pulmonary arteries after the Norwood procedure: comparison between Blalock-Taussig shunt and right ventricular-pulmonary artery conduit. *Ann Thorac Surg* 2008;86(4):1299–304.
- [4] Simsic JM, Bradley SM, Stroud MR, Atz AM. Risk factors for interstage death after the Norwood procedure. *Pediatr Cardiol* 2005;26(4):400–3.
- [5] Ohye RG, Sleeper LA, Mahony L, Newburger JW, Pearson GD, Lu M, et al. Comparison of shunt types in the Norwood procedure for single-ventricle lesions. *N Engl J Med* 2010;362(21):1980–92.